

Agro - techniques of CARI Brinjal 1: A bacterial wilt resistant variety

Sponsored by



**Directorate of
Seed Research**



**P. K. Singh, Krishna Kumar, A.K. Singh, Ajanta Birah, R. K. Gautam, Israr Ahmad
Naresh Kumar, S. K. Zamir Ahmad, K. Sakthivel A.K. Tripathi and S. Dam Roy**

Central Island Agricultural Research Institute

(Indian Council of Agricultural Research)

Post Box No.181, Port Blair – 744 101

(Andaman & Nicobar Islands)



Brinjal (*Solanum melongena* L.) is the most important vegetable crop of Andaman and Nicobar Islands. Its fruits are mainly consumed as cooked vegetable in various ways. It is low in calories fats, protein and fibre but rich in carbohydrates, minerals and vitamins. Bacterial wilt of brinjal caused by *Ralstonia solanacearum* is a serious disease, which limits production from 20 to 50 per cent in Andaman and Nicobar Islands. This disease has risen to alarming proportion in Bay islands and rest of country. Many of the commercial varieties growing are susceptible to this disease due to complexity of pathogen nature and little success has been attained using chemical control measures. The only effective approach of bacterial wilt control is to develop resistant cultivars. Recently the Central Island Agricultural Research Institute, Port Blair has developed a bacterial wilt resistant brinjal variety CARI Brinjal 1 (INGR 12015). Its plants are medium tall, semi spreading type with profuse branching and having light greenish broad leaves. Fruits of this variety are oblong, light green in colour and medium compact with low seediness. CARI Brinjal 1 also exhibited drought tolerant ability during water stress situations and thus suitable for growing in islands conditions during dry season (during October to May).

Climate

It is a warm climate crop and requires a long growing season with the high average day & night temperatures. A daily mean temperature of 15-25°C is most favourable for its production.

Soil and field preparation

Brinjal is moderately deep rooting and can be grown on a wide range of soils types, but well-drained loam soils are much suitable for brinjal crop.

The soil pH should not be > 7.0 for its better growth and development. The cropping period is about 6 months. Therefore, the soils should be well prepared by deep ploughed before transplanting the seedlings. The raised beds should be prepared of 1.25 m breadth for planting.

Sowing Time and Seed Rate

CARI Brinjal 1 can be grown twice in a year. The crop sowing time is May-June and November- December. For one hectare area about 250 to 300 g of seed is required.

Planting and Spacing

For CARI Brinjal 1 optimum spacing is 90 x 90 cm with one plant/hill. Healthy seedlings aged 20 to 25 days with 3-4 fully expanded leaves are suitable for transplanting. After primary and secondary land preparation, level the land and make planting holes at a spacing of 90 x 90 cm in raised beds. Addition of FYM (200 g/hill) to planting holes facilitate quick establishment of the transplant. Transplanting should be done in the evening to avoid mid day heat. Wilted seedling must be light irrigated after transplanting in the beds.

Manures & Fertilizers

Organic manure such as compost or FYM (13-16 t/ha) should be applied 4-5 weeks before transplanting. At the time of land preparation 60 kg urea 150 kg DAP and 80 kg MOP should be applied. After transplanting three doses of Urea @ 75kg/ha must be applied at one month interval as top dressing.

Irrigation

Necessary irrigation is important for getting high fruit yield. Brinjal fields should be regularly irrigated to keep the soil moist during dry days.

Seed treatment

The seed should be treated with Streptocycline (1g/ 40 litres of water) for 30 minutes or Thiram (2g/kg seed) or *Trichoderma* (4-5 g/kg seed) before seed sowing in nursery.

Nursery raising and management

For raising of one heater of brinjal crop 20-21 nursery beds of 3 x 1 m are required. Soil should be prepared well having fine tilth. Prepare raised beds to a height of 15-20cm. Incorporate well decomposed FYM at the rate of 4-5 kg/m². Treated seeds should be sown in beds 10-15 cm apart at the depth of 0.5 to 1 cm and cover with a thin layer of FYM. Spread a layer of straw mulch over beds and water daily. Remove the mulch when the germination is complete. Avoid exposure of seedlings to heavy rains and to prolong sunlight. Seedlings will be transplanting after 20-25 days.

Weed Management

Three to 4 hand weeding at 25-30 days interval are needed for good crop growth and yield. Dark coloured plastic mulches or thatch can also be used but light hand weeding is required to remove weeds that emerge through the planting hole. Chemicals such as Pendimethalin (1.0 kg a.i./ha) or Fluchloralin (1.0 kg a.i. / ha) can be used as pre-emergence herbicide.

Disease management

Damping off, Phomopsis blight and Bacterial wilt are the major diseases that affect this crop. However, CARI Brinjal 1 is resistant to bacterial wilt.

Damping off (*Pythium* sp)

Symptoms of this disease are rotting the base of the nursery plants and then die off. The seedlings are attacked at the collar region and the attacked seedlings are toppled down. The disease spreads through fungi present in the soil. The disease can be controlled by seed treatment with Carbendazim (2g/kg) of seed.

Phomopsis Blight (*Phomopsi vexans*)

This is a common disease in brinjal. The infection starts with foliage blight, but the most destructive phase of the disease is the fruit rot. Pale, sunken spots develop on the fruit and may progress to cover the entire fruit surface. The affected leaves turn yellow and dry. For disease management Foliar spray of Mancozeb 75 % WP (3 g per litre of water) can be applied in the early vegetative stage.

Pest management

Major pests that affect brinjal are Shoot and Fruit borer (SFB) and Jassid.

Shoot and fruit borer (*Leucinodes orbonalis*)

Shoot and fruit borer cause a serious damage to the fruits leading to severe reduction in the yield. The damage to the fruits starts soon after transplanting and continues till harvest of the fruits. Small pinkish larva of the pest initially bore into the terminal shoots resulting in withering and drying of the shoot. In the later stage, it bores into the young fruits by making holes and feeds inside which makes the fruits unfit for consumption. Such fruits rot in severe case. The wilting symptoms first appear in immature shoots soon after observing wilted shoot remove and destroy them. The damage can be controlled by Spinosad 45 SC (0.01 %)

(1ml/10 litre water) or Carbaryl (0.1%) or Rynaxypyr 20% SC @ 40 and 50g a.i. /ha, if sprayed at 11 days interval from fruit setting would give effective control of shoot and fruit borer of brinjal and with appreciably 'no harm to the beneficial natural insect fauna of brinjal.

Jassids (*Amrasca biguttula biguttula*)

Both nymphs and adults suck the sap from the lower surface of the leaves. The infested leaf curl upward along the margins, which may turn yellowish and show, burnt up patches. Fruit setting is adversely affected by the infestation.


Control: Jassids are controlled by spraying Malathion (0.1%) or Dichlorvos (0.05%) 20 days after transplanting.

Beside these the following cultural practices can be used to avoid the pest and diseases.

- ⊙ Healthy seed should be selected for sowing.
- ⊙ Destruction of infected plant material.
- ⊙ Continuous raising of nursery in the same plot should be avoided.
- ⊙ Application of bio-control agent *Trichoderma* in soil (1.5kg/ha).
- ⊙ Removal and destruction of affected plant parts.
- ⊙ Avoid crop rotation with bhindi, tomato.
- ⊙ Seed treatment with *Trichoderma viride* (4-5g/kg) of seed
- ⊙ Soil application with Neem cake @250 kg/ha
- ⊙ Use of yellow sticky traps against White flies
- ⊙ Clipping of shoot borer infested terminals
- ⊙ Leucinodes adult monitoring with pheromone traps

Harvesting & Yield

CARI B-1 fruit are ready for harvest 60–70 days after transplanting or one month after fruit set. Pick fruit in the cooler part of the day to avoid prolonged exposure to the sun. This variety can give yield up to five months. Maximum fruit yield of 30-35 t/ha can be harvested.





Address for correspondence:

DIRECTOR

CENTRAL ISLAND AGRICULTURAL RESEARCH INSTITUTE

Port Blair-744 101, Andaman and Nicobar Islands

Phone- 03192 250341, 250436,

Fax- 03192 251068

Email: directorcaripb@gmail.com